

# FUNIX 25 EC

## 500ml

### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin.

When opening the container and preparing spray wear:

- cotton overalls buttoned to the neck and wrist,
- a washable hat,
- elbow-length butyl rubber gloves and
- face shield or goggles.

When using the prepared spray wear:

- cotton overalls buttoned to the neck and wrist,
- a washable hat and
- elbow-length butyl rubber gloves

If product gets in eyes, wash it out immediately with water.

Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing

### FIRST AID

If poisoning occurs, contact a doctor. If swallowed, DO NOT induce vomiting. Give a glass of water.

### STORAGE

Store in original container in a well-ventilated, cool, dry, and secure area. Ship and store away from food, feed or seed.

### DECONTAMINATION AND DISPOSAL

User should dispose of unwanted active ingredient and containers in accordance with municipal or provincial regulations. For additional details and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

(Difenoconazole 25%EC)

ACTIVE INGREDIENT: Difenoconazole 25%  
Formulation: EC (Emulsifiable Concentrate)

**KEEP OUT OF THE REACH OF CHILDREN  
AND PREVENT ACCESS BY UNAUTHORIZED  
PERSONNEL**

**Mode of action:** Systemic fungicide with preventive and curative action. Absorbed by the leaves, with acropetal and strong translaminar translocation.

**Batch No.:**  
**Date of manufacture:**  
**Expiry Date:**

**CROWNBRIDGE MANAGEMENT L.P.**  
**Suite 260,2323 - 32 Avenue N.8.,**  
**Calgary, Alberta T2E623,**  
**Canada**  
[www.crownbridgetrading.com](http://www.crownbridgetrading.com)

### DIRECTIONS FOR USE

Restraint: DO NOT apply more than 6 applications of this product per season. The effect of FUNIX 25EC could be diminished if rain falls within 2 hours of application.

Crop	Disease	Rate	Critical comments
Asparagus	Stemphylium Leaf Spot	500 ml in 500 to 700 litres of water/ha	After close up of beds, apply at the first sign of the disease and repeat at 3 week intervals with a maximum of 3 applications during active fern growth. The addition of a non-ionic wetting agent at manufacturer's label recommendations is essential to assist coverage. Apply an approved protectant fungicide at other times. <b>Aerial application</b> - apply in a minimum of 100 litres water/ha to achieve satisfactory coverage.
Carrots	Leaf Spots ( <i>Alternaria</i> / <i>Cercospora</i> )		Apply at the first sign of the diseases and repeat at 2 to 3 week intervals with a maximum of 3 applications during active crop growth. Apply an approved protectant fungicide at other times.
Brassicas Horticultural	Ring Spot		Apply at the first signs of the disease and depending on disease pressure, repeat at 2 to 3 week intervals with a maximum of 3 applications during active crop growth. Apply an approved protectant fungicide during crop maturity.
Potatoes	Early Blight		Apply at the first sign of disease and repeat at 2 week intervals with a maximum of 3 applications. Apply an approved protectant fungicide at other times.

### PRECAUTIONS

Re-entry Period: DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist and butyl rubber gloves. Clothing must be laundered after each day's use.

