

MITICIDE ROCKET™ WP

For the control of Two-spotted Mite
and European Red Mite in Apples,
Pears and Peaches and Two-spotted Mite.

ACTIVE INGREDIENT: % BY WT

TEBUFENPYRAD	20.0%
OTHER INGREDIENTS:	80.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Inside for Additional Precautionary Statements and Directions for Use



CROWNBRIDGE

CROWNBRIDGE MANAGEMENT L.P. Suite 260, 2323 - 32 Avenue N.8.
Calgary, Alberta T2E6Z3, Canada www.crownbridgetrading.com

DIRECTIONS FOR USE

Restraints: DO NOT apply if rain is expected within 3 hours.

DONOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Apples, Pears, Peaches	Two-spotted Mite (Tetranychus urticae), European Red Mite (Panonychus ulmi)	Qld, NSW, Vic, Tas, SA, WA, only	Dilute Spraying: 25 to 50 g /100 L Concentrate Spraying: Refer to the Application Section	14 days	Apply by dilute or concentrate spraying equipment. Use higher rate when a longer period of residual control is required. Note: DO NOT use lower rates unless predators are present and IMC is being practiced. Apples and peaches: spray before mite infestation reaches 70% of leaves infested. Pears: spray before mite infestation reaches 25% of leaves infested or 1.4 mites per leaf average. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. When applying with concentrate water volumes, DO NOT use at rates greater than 150 g/100L.
Ornamentals	Two-spotted Mite (Tetranychus urticae)	Qld, NSW, Vic, SA, WA, NT only	5 to 10 g /10 L	-	Lower rate may only suppress two-spotted mite build-up. Application to some varieties of chrysanthemum and gerbera may cause some phototoxicity. Spraying at early signs of mite infestations may extend period of residual control.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

Apples, Pears, Peaches:

Ornamentals:

DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

NOT REQUIRED WHEN USED AS DIRECTED.

DO NOT GRAZE ANY TREATED AREAS.

DO NOT FEED PRODUCE FROM TREATED AREAS TO ANIMALS, INCLUDING POULTRY.

GENERAL INSTRUCTIONS

MIXING

Add the required quantity of product to clean water while stirring or while agitators are in motion. Maintain agitation during spraying. Note: DO NOT open the water-soluble sachets.

APPLICATION

The product may be applied as a dilute or concentrate spray.

DILUTE SPRAYING (Apples, Pears, Peaches only):

Use a sprayer designed to apply high volumes of water up to the point of run off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run off. Avoid excessive run off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

CONCENTRATE SPRAYING (Apples, Pears, Peaches only):

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only:

Dilute spray volume as determined above: For example, 1500 L/ha.

Your chosen concentrate spray volume: For example, 500 L/ha.

The concentration factor in this example is: 3 X (ie 1500 L (500 L = 3)

If the dilute label rate is 25 g / 100L, then the concentrate rate becomes 3 x 25, that is 75 g/100 L of concentrate spray. The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and following industry Best Practices.

Note: in some instances, residual control may be reduced from the use of concentrate sprays.

COMPATIBILITY

A number of insecticides, fungicides, spreaders, adhesives and wetting agents are compatible with ROCKET if mixed in the spray tank and used promptly. As it is impossible to know all the commercial products on sale, we cannot assume any responsibility for tank mix compatibility. We can only recommend to the user that a preliminary compatibility test be carried out with the planned mixture to observe the physical aspect of the spray and the reaction it may produce on the plants to be treated during the days following the test.

RESISTANCE MANAGEMENT

Apples/Pears/Peaches: A successful resistance management strategy will use a variety of methods including: use of chemicals from different groups (e.g. oils, organophosphates) and use of predators.

Make only one application of ROCKET per season. If another spray is required, alternate with products from other chemical groups. If ROCKET is the last miticide applied one season then the first miticide spray the following season should be with a product from another chemical group.

Ornamentals: A successful resistance management strategy will use a variety of methods including: use of chemicals from different groups (e.g. oils, organophosphates) and use of predators. Use no more than 4 sprays of ROCKET each calendar year. DO NOT apply ROCKET sequentially. Preferably use at least 3 miticide sprays from 2 other chemical groups before re-using ROCKET.

Integrated Mite Control (IMC):

ROCKET, when applied within the labelled rate ranges, allows IMC (based on *T. pyri* and *T. occidentalis*) to be maintained - but will adversely affect the mite predator *Phytoseiulus persimilis*.

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management ROCKET is a Group 21A insecticide.

GROUP	21A	INSECTICIDE
-------	------------	-------------

Some naturally occurring insect biotypes resistant to ROCKET and other Group 21A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if ROCKET and other Group 21A insecticides are used repeatedly. The effectiveness of ROCKET on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Crownbridge Management L.P accepts no liability for any losses that may result from failure of ROCKET to control resistant insects. ROCKET may be subject to specific resistance management strategies. For more information contact your supplier, Crownbridge Management L.P representative or local agricultural department agronomist.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, ENVIRONMENT AND BEES

Dangerous to fish and aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT spray across open bodies of water. Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. May adversely affect the mite predator *Phytoseiulus persimilis*.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes. Avoid inhaling dust. When preparing spray wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash contaminated clothing.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.crownbridgetrading.com

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Canadian Consumer Law, Crownbridge Management L.P. liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Crownbridge Management L.P. option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Crownbridge Management L.P. or the place in which you purchased this product for a full refund.

Batch No:

Date of Manufacture:

G.H.S. STATEMENTS

HAZARDS H312: Harmful in contact with skin. H332: Harmful if inhaled.

PREVENTION P262: Do not get in eyes, on skin, or on clothing. P270: Do not eat, drink or smoke when using this product. P271: Use only outdoors or in a well-ventilated area. P280: Wear protective gloves, protective clothing and eye or face protection.

RESPONSE P361: Remove all contaminated clothing immediately. P301+P330+P331: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. P302+P352: IF ON SKIN: Wash with plenty of soap and water. P304+P340: IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. P370+P378: In case of fire, note the following. Water fog or fine spray is the preferred medium for large fires. Try to contain spills, minimise spillage entering drains or water courses.